


---

## Frequency of Screening Mammography Study



---

A study that compared the benefits and harms of the frequency of screening mammography to age, breast density and postmenopausal use of hormone therapy (HT) suggests that women ages 50 to 74 years who undergo biennial screenings have a similar risk of advanced-stage disease and a lower cumulative risk of false-positive results than those who get mammograms annually, according to a report published Online First by JAMA Internal Medicine, a JAMA Network publication.

The U.S. Preventive Services Task Force 2009 guidelines recommended biennial mammography, rather than the previously recommended mammography every one to two years, for women ages 50 to 74, but the updated guidelines did not consider the influence of breast cancer risk factors beyond age, according to the study background.

Karla Kerlikowske, M.D., of the University of California, San Francisco, and colleagues sought to determine whether the benefits, such as early detection, and the harms, such as a false-positive mammography result or biopsy recommendation, differ among women undergoing screening mammography according to age, breast density and postmenopausal HT use.

Researchers analysed data collected from January 1994 to December 2008 from mammography facilities in community practice that participate in the Breast Cancer Surveillance Consortium (BCSC) mammography registries. Data were collected from 11,474 women with breast cancer and 922,624 without breast cancer.

The authors found that biennial vs. annual mammography for women ages 50 to 74 was not associated with an increased risk of advanced-stage or large-size tumours regardless of a woman's breast density or HT use. However, the results indicate that among women ages 40 to 49 years with extremely dense breasts, biennial mammography vs. annual was associated with an increased risk of advanced-stage cancer (odds ratio [OR], 1.89) and large tumours (OR, 2.39).

Study results also show that the cumulative probability of a false-positive mammography result was high among women undergoing annual mammography with extremely dense breasts who were either ages 40 to 49 years (65.5 percent) or used oestrogen plus progestogen (65.8 percent) and was lower among women ages 50 to 74 years who underwent biennial or triennial mammography with scattered fibroglandular densities or fatty breasts.

The study concludes, "Women aged 50 to 74 years, regardless of breast density or HT use, can undergo biennial rather than annual mammography because biennial screening does not increase the risk of presenting with advanced disease but does substantially reduce the cumulative risk of a false-positive mammography result and biopsy recommendation. Women aged 40 to 49 years with extremely dense breasts who choose to undergo mammography should consider annual screening to decrease the risk of advanced-stage disease but should be informed that annual screening leads to a high cumulative probability of a false-positive mammography result because of the additional screening examinations."

JAMA Intern Med. Published online March 18, 2013. doi:10.1001/jamainternmed.2013.307.

Published on : Tue, 19 Mar 2013